


Future prospects for charm physics at BESIII and beyond



BESIII

Yangheng Zheng
on behalf of BESIII collaboration
University of Chinese Academy of Sciences


May. 22, 2015

Outline

- ◆ **Current BEPCII/BESIII**
 - ◆ **Operations and data taking plans**
- ◆ **Detector upgrade**
 - ◆ **Endcap TOF**
 - ◆ **Inner tracking detector**
- ◆ **Prospects for charm physics**
 - ◆ **BESIII**
 - ◆ **Super Tau-charm factory**
- ◆ **Summary**


Current status of BEPCII

◆ BEPCII: A double ring e^+e^- collider



- ◆ A record $L_{\text{peak}} = 8.5 \times 10^{32} \text{ cm}^{-2}\text{s}^{-1}$ @ 3.773 GeV reached in Nov. 20, 2014
- ◆ Weekly record luminosity: 169 pb^{-1} , Average of daily lumi: $>17 \text{ pb}^{-1}$
- ◆ 500 pb^{-1} data collected @ 4.6 GeV : the limit of BEPCII energy region
- ◆ Bunch crossing time: $8 \text{ ns} \rightarrow 6 \text{ ns}$


Data samples available



- ◆ 20 points for R & QCD Scan: 500 pb^{-1} finished in May 1st, 2015
- ◆ Currently: at $Y(2175)$ resonance, plan: 100 pb^{-1}

Data samples @ charm threshold

- ◆ CLEO-c: 818 pb⁻¹ @ $\psi(3770)$
- ◆ BESIII: 2.9 fb⁻¹ (~3.5 x CLEO-c data) @ $\psi(3770)$



- ◆ BESIII: 0.5 fb⁻¹ @ $\psi(4040)$
- ◆ In 2015-2016 run period: 3 fb⁻¹ @ 4.17 GeV

Data taking plan



Approved plan for 2015/2016 run period

- ◆ 4170 data taking 3 fb⁻¹, **5 months**
- ◆ Data taking at around χ_{c1} mass **25 days**
 - ◆ e⁺e⁻ → χ_{c1} , Collins fragmentation function
 - 2-3 points, 3.5136 (3 MeV below, 20 pb⁻¹)
 - 3.5106 (180 pb⁻¹)
 - 3.5100 (0.5 MeV below) or χ_{c1} mass for 180 pb⁻¹
- ◆ Psi' scan 500 pb⁻¹ **25 days**

Final goal for data set at threshold: ~20 fb⁻¹

BESIII detector upgrade

- ◆ Inner Drift Chamber appeared aging effect
 - ◆ Cathode aging → Malter discharge
 - ◆ Adding ~2000ppm water vapor into the gases
 - ◆ Anode aging → gain drop 14% - 26%
 - ◆ Short-term upgrade plan: **a new inner drift chamber**
 - ◆ Long-term upgrade plan: **a 3-layer CGEMs inner tracker**
- ◆ Endcap TOF (ETO: scintillator + PMT)
 - ◆ To improve PID
 - ◆ Upgrade with MRPC
 - ◆ Less affected by scattering
 - ◆ Tracking with more readout pads
 - ◆ Total resolution: ~140ps → <80ps



Upgrade for inner tracking detector




◆ New inner drift chamber

- ◆ Chamber wiring finished in January, 2015
- ◆ Ready for installation





◆ CGEM inner tracker

- ◆ Proposed by the Italy group;
- ◆ 3-Layer CGEM foils (KLOE2-like);
- ◆ The design is coming to a conclusion;
- ◆ construction of the layer 2 has started;
- ◆ beam test of the prototype is ongoing;
- ◆ software: progress in simulation and reconstruction.




Endcap TOF upgrade



- ◆ **Two MRPCs has been installed.**
 - ◆ Testing with real data
 - ◆ A VERY preliminary calibration \Rightarrow time resolution ~ 70 ps
- ◆ **The production of the whole ETOF system: finished**
- ◆ **Performance and stability test is carrying**
 - ◆ on ~ 4 months cosmic-ray test
- ◆ **Simulation: new GDML and Digitization is developed.**

Physics at tau-charm Energy Region



- Hadron form factors
- $Y(2175)$ resonance
- Multiquark states with s quark, Zs
- MLLA/LPHD and QCD sum rule predictions

- Light hadron spectroscopy
- Gluonic and exotic states
- Process of LFV and CPV
- Rare and forbidden decays
- Physics with τ lepton

- XYZ particles
- D mesons
- f_D and f_{D_s}
- D_0 - \bar{D}_0 mixing
- Charm baryons

Charm facilities

- ◆ **Hadron colliders (huge cross-section, energy boost)**
 - ◆ Tevetron (CDF, D0)
 - ◆ LHC (LHCb, CMS, ATLAS)
- ◆ **e^+e^- Colliders (more kinematic constrains, clean environment, ~100% trigger efficiency)**
 - ◆ B-factories (Belle, BaBar)
 - ◆ **Threshold production (CLEOc, BESIII)**
 - ◆ **Can not compete in statistics with Hadron colliders & B-factories! ! !**
 - ◆ Quantum Correlations (QC) and CP-tagging are unique
 - ◆ Only D meson pairs, no extra CM Energy for pions
 - ◆ Systematic uncertainties cancellations while applying double tag technique

Physics at Charm threshold


- ◆ Decay constants & form factors for Charm meson
- ◆ Quantum correlations at $\psi(3770)$
 - ◆ CPV measurements
 - ◆ Strong phase measurements
- ◆ Rare decays
- ◆ Charm baryons
- ◆ D^0 - \underline{D}^0 mixing & CPV @ $\psi(4040)$

Many new BESIII results have been released!

Selected results will be shown!

Double Tag (DT) techniques

- ◆ 100% of beam energy converted to D pair (Clean environment, kinematic constrains ν Recon.)
- ◆ D generated in pair \Rightarrow absolute Branching fractions
- ◆ At $\psi(3770)$ charm production is $D^0\bar{D}^0$ and D^+D^-
- ◆ Fully reconstruct about 15% of D decays




$$\Delta E = E_D - E_{\text{Beam}}$$

$$M_{\text{BC}} = \sqrt{E_{\text{Beam}}^2 - p_D^2}$$

- ◆ Double tag techniques: Hadronic tag on one side, on the other side for leptonic/semileptonic studies. Neutrino is reconstructed from missing energy and momentum (Double tag efficiency is high.)

$f_{D(s)^+}$: Leptonic decays




The Feynman diagram shows a charm quark (c) and an anti-down quark (\bar{d}) forming a D^+ meson. They interact via a W^+ boson, which then decays into a lepton (ℓ^+) and a neutrino (ν).

$$\Gamma(D^+ \rightarrow \ell^+ \nu_\ell) = f_D^2 |V_{cd}|^2 \frac{G_F^2}{8\pi} m_D m_\ell^2 \left(1 - \frac{m_\ell^2}{m_D^2}\right)^2$$

- ◆ Extract decay constant $f_{D(s)}$ incorporates the strong interaction effects (wave function at the origin)
 - ◆ Multiple tests with charm: f_D , f_{D_s} and f_D/f_{D_s}
- ◆ To validate Lattice QCD calculation of $f_{B(s)}$ and provide constrain of CKM-unitarity
- ◆ Sensitive to New Physics (Charged Higgs contribution, ...)

f_{D^+} Results ($D^+ \rightarrow \mu^+ \nu$)



Phys. Rev. D89, 051104

◆ 2.92fb^{-1} @ 3.773 GeV

◆ Muon counter information applied

◆ Kinematic variable: M_{miss}^2

$$M_{\text{miss}}^2 = (E_{\text{Beam}} - E_{\mu})^2 - (-\vec{p}_{\text{tag}} - \vec{p}_{\mu})^2 \geq 0$$

◆ 451 $D \rightarrow \mu \nu$ candidates observed

◆ Low background

$$N(D^+ \rightarrow \mu^+ \nu) = 409.0 \pm 21.2 \pm 2.3$$

$$B(D^+ \rightarrow \mu^+ \nu) = (3.71 \pm 0.19 \pm 0.06) \times 10^{-4}$$


$$f_{D^+} = (203.2 \pm 5.3 \pm 1.8) \leftarrow |V_{cd}| \text{ of CKM-Fitter Input}$$

$$|V_{cd}| = 0.221 \pm 0.006 \pm 0.005$$


\leftarrow LQCD calculated $f_D = 207 \pm 4 \text{ MeV}$ [PRL100(2008)062002]

The error is still dominated by statistics! More data is needed.

$f_{D(s)^+}$ Comparison



BESIII: 2.7% with 2.92fb^{-1}
 BESIII final: 1.5% with 10fb^{-1}



CLEO-c: 2.5% with 0.68fb^{-1}
 BESIII final: 1.25% with 5fb^{-1}

Form Factors: Semileptonic decays

The diagram shows a D meson (quarks c and \bar{q}) decaying into a P meson (quarks q' and \bar{q}) through a W^+ boson. The W^+ boson then decays into a lepton pair (ν_e and e^+).

$$\frac{d\Gamma(D \rightarrow K(\pi) e \nu)}{dq^2} = \frac{G_F^2 |V_{cs(d)}|^2 P_{K(\pi)}^3}{24 \pi^3} |f_+(q^2)|^2$$

$q^2 = (p_l + p_\nu)^2 \Rightarrow M_{\text{inv}}^2$
of lepton pair

◆ $D_{(s)} \rightarrow P l \nu$ (Theoretically clean)

◆ Measure $|V_{cx}|$ x FF

◆ Charm physics:



◆ CKM-unitarity $\Rightarrow |V_{cx}|$, extract FF, test LQCD

◆ Input LQCD FF to test CKM-unitarity

◆ B physics: Validate LQCD for form factor, extract $|V_{ub}|$ to test CKM-unitarity

◆ Example: $B \rightarrow \pi l \nu \Rightarrow |V_{ub}| = 3.92 \pm 0.09 \pm 0.45$ (Theory) rely on LQCD Form Factor calculations (provide perfect calibration)

Form Factors fit results ($D^0 \rightarrow K/\pi e^+ \nu$)



	$D^0 \rightarrow K e^+ \nu$		$D^0 \rightarrow \pi e^+ \nu$	
Simple Pole	$f_K^+(0) V_{cs} $	$0.7209 \pm 0.0022 \pm 0.0033$	$f_\pi^+(0) V_{cd} $	$0.1475 \pm 0.0014 \pm 0.0005$
	M_{pole}	$1.9207 \pm 0.0103 \pm 0.0069$	M_{pole}	$1.9114 \pm 0.0118 \pm 0.0038$
Mod. Pole	$f_K^+(0) V_{cs} $	$0.7163 \pm 0.0024 \pm 0.0034$	$f_\pi^+(0) V_{cd} $	$0.1437 \pm 0.0017 \pm 0.0008$
	α	$0.3088 \pm 0.0195 \pm 0.0129$	α	$0.2794 \pm 0.0345 \pm 0.0113$
ISGW2	$f_K^+(0) V_{cs} $	$0.7139 \pm 0.0023 \pm 0.0034$	$f_\pi^+(0) V_{cd} $	$0.1415 \pm 0.0016 \pm 0.0006$
	r_{ISGW2}	$1.6000 \pm 0.0141 \pm 0.0091$	r_{ISGW2}	$2.0688 \pm 0.0394 \pm 0.0124$
Series.2.Par	$f_K^+(0) V_{cs} $	$0.7172 \pm 0.0025 \pm 0.0035$	$f_\pi^+(0) V_{cd} $	$0.1435 \pm 0.0018 \pm 0.0009$
	r_1	$-2.2278 \pm 0.0864 \pm 0.0575$	r_1	$-2.0365 \pm 0.0807 \pm 0.0260$
Series.3.Par	$f_K^+(0) V_{cs} $	$0.7196 \pm 0.0035 \pm 0.0041$	$f_\pi^+(0) V_{cd} $	$0.1420 \pm 0.0024 \pm 0.0010$
	r_1	$-2.3331 \pm 0.1587 \pm 0.0804$	r_1	$-1.8434 \pm 0.2212 \pm 0.0690$
	r_2	$3.4223 \pm 3.9090 \pm 2.4092$	r_2	$-1.3871 \pm 1.4615 \pm 0.4677$

Comparison of Form Factors

Gang Rong
CKM2014

Theory:

HPQCD (2010) $0.747 \pm 0.011 \pm 0.015$

Fermilab/MILC (2005) $0.73 \pm 0.03 \pm 0.07$

Sum Rules (2009) $0.75^{+0.11}_{-0.08}$

Experiment:



CLEO-c (2009) $0.739 \pm 0.007 \pm 0.005 \pm 0.000$

Belle (2006) $0.695 \pm 0.007 \pm 0.022$

BaBar (2007) $0.727 \pm 0.007 \pm 0.005 \pm 0.007$

BESIII (2014) $0.7368 \pm 0.0026 \pm 0.0036$

Based on preliminary result





BESIII: the most precise measurements

The error of $f_+^{D \rightarrow \pi}$ is still dominated by statistics.


CKM matrix elements $|V_{cd}(s)|$

Gang Rong
CKM2014




BESIII: the most precise measurements

Reaches for rare charm decays?



Rare decays ($D^0 \rightarrow \gamma\gamma$)


arXiv:1505.03087



Double tag method

Tag modes

$$\left. \begin{aligned} \bar{D}^0 &\rightarrow K\pi \\ \bar{D}^0 &\rightarrow K\pi\pi^0 \\ \bar{D}^0 &\rightarrow K3\pi \\ \bar{D}^0 &\rightarrow K3\pi\pi^0 \\ \bar{D}^0 &\rightarrow K\pi2\pi^0 \end{aligned} \right\} D^0 \rightarrow \gamma\gamma$$




Best limit

$$\mathcal{B}_{D^0 \rightarrow \gamma\gamma} < 2.2 \cdot 10^{-6} \text{ @ } 90\% \text{ C.L.}$$



BaBar 470 fb^{-1} @ $\sqrt{s}(4S)$
PRD85,091107(2012)

$$\mathcal{B}(D^0 \rightarrow \pi^0\pi^0) = (8.24 \pm 0.21(\text{stat.}) \pm 0.30(\text{syst.})) \times 10^{-4}$$



BESIII 2.92 fb^{-1} @ 3770:

$$\mathcal{B}(D^0 \rightarrow \gamma\gamma) < 3.8 \times 10^{-6} \text{ @ } 90\% \text{ C.L.}$$


BESIII 10 fb^{-1} @ 3770:

$$\mathcal{B}_{D^0 \rightarrow \gamma\gamma} < 1.0 \times 10^{-6}$$

BESIII has much smaller background than that at B factory, peaking background from $D^0 \rightarrow \pi^0\pi^0$ is under control.

$D^0 \rightarrow \gamma\gamma$ reach at super τ -charm

1 ab⁻¹ at threshold
at **super τ -charm**
factory will reach
Long Distance
contribution:
about 60 events are
expected per year.



δ and γ/ϕ_3 input

- ◆ D hadronic parameters for a final state

$$f: \frac{A(\bar{D}^0 \rightarrow f)}{A(D^0 \rightarrow f)} \equiv -r_D e^{-i\delta_D}$$

- ◆ Charm mixing parameters: $x = \frac{\Delta M}{\Gamma}$, $y = \frac{\Delta\Gamma}{2\Gamma}$

- ◆ Time-dependent WS $D^0 \rightarrow K^+ \pi^-$ rate \Rightarrow

$$y' = y \cos \delta_{K\pi} - x \sin \delta_{K\pi} = (0.72 \pm 0.24)\% \text{ (LHCb 2012)}$$

- ◆ $\delta_{K\pi}$: QC measurements from Charm factory

- ◆ γ/ϕ_3 measurements from $B \rightarrow D^0 K$

- ◆ $b \rightarrow u$: $\gamma/\phi_3 = \arg V_{ub}^*$

- ◆ most sensitive method to constrain γ/ϕ_3 at present

- ◆ GLW, ADS method

- ◆ r_D, δ_D : QC measurements from Charm factory

- ◆ GGSZ method

- ◆ c_i, s_i : QC measurements from Charm factory

Time-integrated decay rates

◆ No time dependent information at Charm threshold

◆ Anti-symmetric wavefunction:

$$\Gamma_{ij}^2 = |\langle i|D^0\rangle\langle j|\bar{D}^0\rangle - \langle j|D^0\rangle\langle i|\bar{D}^0\rangle|^2$$

◆ Double tag rates:

$$A_i^2 A_j^2 [1 + r_i^2 r_j^2 - 2r_i r_j \cos(\delta_i + \delta_j)]$$

◆ CP tag: $r=1, \delta=0$ or π ; l^\pm tag: $r=0$

◆ Single and Double tag rates

$$\text{◆ } z_f \equiv 2 \cos \delta_f, r_f \equiv \frac{A_{DCS}}{A_{CF}}, R_M \approx \frac{x^2 + y^2}{2}$$

$$\psi(3770) \rightarrow [D^0 \bar{D}^0 - \bar{D}^0 D^0] / \sqrt{2}$$

$$= -[D_{CP+} D_{CP-} - D_{CP-} D_{CP+}] / \sqrt{2}$$

$$D_{CP\pm} = [D^0 \pm \bar{D}^0] / \sqrt{2}$$


C-odd	f	\bar{f}	l^+	l^-	CP+	CP-
f	$R_M [1 + r_f^2 (2 - z_f^2) + r_f^4]$					
\bar{f}	$1 + r_f^2 (2 - z_f^2) + r_f^4$	$R_M [1 + r_f^2 (2 - z_f^2) + r_f^4]$				
l^+	r_f^2	1	R_M			
l^-	1	r_f^2	1	R_M		
CP+	$1 + r_f (r_f + z_f)$	$1 + r_f (r_f + z_f)$	1	1	0	
CP-	$1 + r_f (r_f - z_f)$	$1 + r_f (r_f - z_f)$	1	1	4	0
Single Tag	$1 + r_f^2 - r_f z_f (A - y)$		1	$2[1 \pm (A - y)]$		

$\delta_{K\pi}$ in $D \rightarrow K\pi$ (BESIII: 2.9 fb^{-1})

PLB 734, 227(2014)

A simple picture: $\frac{\langle K\pi | \overline{D}^0 \rangle}{\langle K\pi | D^0 \rangle} \equiv \frac{\overline{A_{K\pi}}}{A_{K\pi}} \equiv r_{K\pi} e^{i\delta_{K\pi}}$

$$\langle K\pi | D_{CP\pm} \rangle = (\langle K\pi | D^0 \rangle \pm \langle K\pi | \overline{D}^0 \rangle) / \sqrt{2} \Rightarrow \sqrt{2} A_{CP\pm} = A_{K\pi} \pm \overline{A_{K\pi}}$$



◆ Measuring $\delta_{K\pi}$ from rate differences if using external $r_{K\pi}$

◆ Reconstructed modes:

◆ Flavor tags: $K^-\pi^+$, $K^+\pi^-$


◆ CP+ tags (5 modes): K^-K^+ , $\pi^+\pi^-$, $K_S^0\pi^0\pi^0$, $\pi^0\pi^0$, $\rho^0\pi^0$

◆ CP- tags (3 modes): $K_S^0\pi^0$, $K_S^0\eta$, $K_S^0\omega$


$\delta_{K\pi}$ in $D \rightarrow K\pi$ (BESIII: 2.9 fb^{-1})

PLB 734, 227(2014)

Single Tags



Double Tags



Direct result:

$$A_{CP \rightarrow K\pi} = (12.7 \pm 1.3(\text{Stat.}) \pm 0.7(\text{sys.}))\%$$

$$2r_{K\pi} \cos \delta_{K\pi} + y = (1 + R_{WS}) \cdot A_{CP \rightarrow K\pi}$$

Using external input for $r_{K\pi}^2$, y , R_{WS} we extract:

$$\cos \delta_{K\pi} = 1.02 \pm 0.11 \pm 0.06 \pm 0.01$$


If BESIII accumulate 10 fb^{-1} on threshold D data:

sensitivity of $\cos \delta_{K\pi} \sim 0.06$

y_{CP} measurement (BESIII: 2.9 fb^{-1})

PLB 744, 339(2015)

We measure the y_{CP} using CP -tagged semi-leptonic D decays, which allows to access CP asymmetry in mixing and decays.



◆ Single Tag decay rate (CP tags)

$$\text{◆ } \Gamma_{CP\pm} \propto 2|A_{CP\pm}|^2(1 \mp y)$$

◆ Double Tag decay rate (Flavor tags + CP tags)

$$\text{◆ } \Gamma_{l;CP\pm} \propto |A_l|^2|A_{CP\pm}|^2$$

◆ Neglect term y^2 or higher order

$$\text{◆ } y_{CP} \approx \frac{1}{4} \left(\frac{\Gamma_{l;CP+}\Gamma_{CP-}}{\Gamma_{l;CP-}\Gamma_{CP+}} - \frac{\Gamma_{l;CP-}\Gamma_{CP+}}{\Gamma_{l;CP+}\Gamma_{CP-}} \right)$$

◆ Reconstructed modes:

◆ Flavor tags: $Ke\nu_e, K\mu\nu_\mu$


◆ CP+ tags (3 modes): $K^-K^+, \pi^+\pi^-, K_S^0\pi^0\pi^0$

◆ CP- tags (3 modes): $K_S^0\pi^0, K_S^0\eta, K_S^0\omega$


y_{CP} measurement (BESIII: 2.9 fb⁻¹)

PLB 744, 339(2015)


Single Tags



Double Tags




- ◆ **BESIII results:**
 $y_{CP} = (-2.0 \pm 1.3 \pm 0.7)\%$
- ◆ **Result is statistically limited**
- ◆ **Systematic uncertainty is relative small**
- ◆ **Most precise measurement with QC charm mesons**
- ◆ **In the limit of no CP violation:**
 $y_{CP} = \mathcal{Y}$
- ◆ **Super τ -C: $\Delta(y_{CP}) \sim 0.1\%$**



$K_S \pi^+ \pi^-$ (BESIII preliminary: 2.9 fb^{-1})

◆ Extract c_i, s_i for “ γ/ϕ_3 GGSZ method”

◆ Preliminary results presented @ APS meeting, Apr. 2014



Bins	c_i		s_i	
	BES-III	CLEO-c	BES-III	CLEO-c
1	0.066 ± 0.066	-0.009 ± 0.088	-0.843 ± 0.119	-0.438 ± 0.184
2	0.796 ± 0.061	0.900 ± 0.106	-0.357 ± 0.148	-0.490 ± 0.295
3	0.361 ± 0.125	0.292 ± 0.168	-0.962 ± 0.258	-1.243 ± 0.341
4	-0.985 ± 0.017	-0.890 ± 0.041	-0.090 ± 0.093	-0.119 ± 0.141
5	-0.278 ± 0.056	-0.208 ± 0.085	0.778 ± 0.092	0.853 ± 0.123
6	0.267 ± 0.119	0.258 ± 0.155	0.635 ± 0.293	0.984 ± 0.357
7	0.902 ± 0.017	0.869 ± 0.034	-0.018 ± 0.103	-0.041 ± 0.132
8	0.888 ± 0.036	0.798 ± 0.070	-0.301 ± 0.140	-0.107 ± 0.240

***Only statistical uncertainty is listed

□ Model prediction
● BESIII
▼ CLEO-c

BESIII Preliminary

Consistent agreement with CLEO-c measurements.


Source: CLEO Collaboration, Physical Review D, vol 82., pp. 112006 - 112035

Charm baryon Λ_c^\pm decays



- ◆ BESIII: Λ_c^\pm Pair production at threshold (4.6GeV)
- ◆ Largest data set @4.6 GeV
- ◆ Double Tag \Rightarrow Model-independent absolute Λ_c^\pm decay Branching fractions

Belle: first model-independent Measurement (PRL113,042002)

ST Λ_c^\pm yields




DT Λ_c^\pm yields



Decay modes	global fit \mathcal{B}	PDG \mathcal{B}	Belle \mathcal{B}
pK_S	1.48 ± 0.08	1.15 ± 0.30	$6.84 \pm 0.24^{+0.21}_{-0.27}$
$pK^- \pi^+$	5.77 ± 0.27	5.0 ± 1.3	
$pK_S \pi^0$	1.77 ± 0.12	1.65 ± 0.50	
$pK_S \pi^+ \pi^-$	1.43 ± 0.10	1.30 ± 0.35	
$pK^- \pi^+ \pi^0$	4.25 ± 0.22	3.4 ± 1.0	
$\Lambda \pi^+$	1.20 ± 0.07	1.07 ± 0.28	
$\Lambda \pi^+ \pi^0$	6.70 ± 0.35	3.6 ± 1.3	
$\Lambda \pi^+ \pi^- \pi^+$	3.67 ± 0.23	2.6 ± 0.7	
$\Sigma^0 \pi^+$	1.28 ± 0.08	1.05 ± 0.28	
$\Sigma^+ \pi^0$	1.18 ± 0.11	1.00 ± 0.34	
$\Sigma^+ \pi^+ \pi^-$	3.58 ± 0.22	3.6 ± 1.0	
$\Sigma^+ \omega$	1.47 ± 0.18	2.7 ± 1.0	

Absolute BR for $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$

- ◆ Dominated process: $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$
- ◆ Input for LQCD calculations
- ◆ First direct absolute BF measurement
- ◆ Theoretical predictions: 1.4% ~ 9.2%



$$B(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e) = (3.63 \pm 0.38 \pm 0.??)\%$$

Statistical error only!

Statistical limited measurement!


Prospects for Λ_c decays

Can BEPCII challenge the CM energy limit at **4.6 GeV**?

With larger Λ_c data sample

- ◆ PWA \Rightarrow intermediate structures in 3-body decays
- ◆ More semileptonic decays: $n l \nu$, $\Lambda^* l \nu$, $\Sigma X l \nu$...
- ◆ Decay asymmetry parameters $\alpha \Leftarrow \Lambda_c^+ \rightarrow BP/BV$
- ◆ Λ_c^+ Rare decays search
 - ◆ Weak radiative decay $\Lambda_c^+ \rightarrow \gamma \Sigma^+$
 - ◆ FCNC $\Lambda_c^+ \rightarrow p l^+ l^-$
 - ◆ LNV $\Lambda_c^+ \rightarrow p e \mu$

Current dataset @4.6 GeV




How about @peak 4.63 GeV?

FIG. 4: The cross section for the exclusive process $e^+e^- \rightarrow \Lambda_c^+ l^-$

Future charm facilities (CHARM 2013)

Novosibirsk machine




Future charm facilities

Marcello A. Giorgi

INFN and Università di Pisa

CHARM 2013- Manchester 31 August-4 September , 2013

The Italian Tau-Charm




Another proposed Machine

- ◆ In China
- ◆ e+e- collider
- ◆ Wide c.m. energy coverage: **2 – 7 GeV**
- ◆ Collider + 4th generation SR source
 - ◆ Symmetric two ring collision
 - ◆ Collision & SR: sharing mode feasible
- ◆ Peak luminosity: **$1 \times 10^{35} \text{ cm}^{-2}\text{s}^{-1}$** (Optimized @ $E_{\text{cm}} = 4 \text{ GeV}$)
- ◆ Data set: **1 ab^{-1}** at Charm threshold
- ◆ Polarized beam
 - ◆ Polarized electron beam source
 - ◆ Siberian Snake curing depolarization

High Intensity Electron Positron Accelerator

HIEPA project

$E_{cm} = 2-7 \text{ GeV}$, $L = 1 \times 10^{35} \text{ cm}^{-2}\text{s}^{-1}$ at 4 GeV




A potential location: Hefei



- ◆ **University of Science and Technology of China (USTC)**
- ◆ **National Synchrotron Radiation Lab and Hefei Light Source, operated by USTC**
- ◆ **The only National Lab operated by University in China. (Totally Four National Labs in China)**

Detector



MUD

- μ/π suppression power >10/30

EMC

- Energy range: 0.02-2.5 GeV
- At 1 GeV σ_E (%)
 - Barrel(CsI): 2
 - Endcap (Cs): 4

PID

- π/K (and K/p) 3-4 σ separation up to 2GeV/c

MDC

- σ_{xy} =130 mm
- $dE/dx < 7\%$, $\sigma_p/p = 0.5\%$ at 1 GeV

PXD

- Material budget $\sim 0.15\% X_0/\text{layer}$
- σ_{xy} =50 mm

Expected Key features

- ◆ **Vertex** – very low material budget $\sim 0.15\text{-}0.3\%X_0/\text{layer}$, $<50\mu\text{m}$ position resolution;
- ◆ **MDC** – pT resolution @ $1\text{GeV}/c$ $0.5\sim 0.7\%$, dE/dx resolution $<7\%$, low material budget ;
- ◆ **PID** – π/K (and K/p) $3\text{-}4\sigma$ separation up to $2\text{GeV}/c$, low material ($<0.5X_0$);
- ◆ **EMC** – stochastic term $<2\%/\sqrt{E}$, constant term $<0.75\%$;
- ◆ **MUD** - μ/π suppression power >10 .

Some sensitivities at HIEPA

- ◆ **With 1 ab^{-1} data at threshold**
 - ◆ **Direct CP violation in $D^+ \rightarrow hh$ sensitivity: $10^{-3} \sim 10^{-4}$**
 - ◆ **Probe y : $\Delta(y_{\text{CP}}) \sim 0.1\%$**
 - ◆ **RM = $(x^2 + y^2)/2 \sim 10^{-5}$ in $K\pi$ and $K\eta$ channels**
 - ◆ **$\Delta(\cos\delta_{K\pi}) \sim 0.007$; $\Delta(\delta_{K\pi}) \sim 2^\circ$**
- ◆ **Clean background and better systematic control in threshold production (complementary to the future B factory results)**

HIEPA related activities

- ◆ **Several domestic workshops**
- ◆ **Jan 13-16, 2015, HIEPA International Workshop on Physics at Future High Intensity Collider @ 2 – 7 GeV in Hefei, China**
- ◆ **June 3 – 4, 2015, Domestic workshop on “Physics, applications and Key technologies on 2 – 7 GeV HIEPA”,**
 - ◆ **more “official” discussions within HEP community in China**
- ◆ **CDR for accelerator & detector in progress (Will be ready by 2016)**

Summary

- ◆ Many BESIII Charm results are released in this conference! It's just the beginning!
- ◆ Charm at threshold provides opportunities for both QCD and New Physics
 - ◆ Very active on XYZ analyses
 - ◆ Will provide best measurements: $f_{D(s)}$ & FF
 - ◆ Unique access to strong phases & ability to extract model-independent results with charm at threshold
 - ◆ Charm baryon results
- ◆ BESIII team has learned and developed technology with charm at threshold.
- ◆ BESIII will continue to run 6 – 8 years.
- ◆ It is natural to propose the e^+e^- intensity frontier for the τ -charm energy region in China \Rightarrow **High Intensity Electron Positron Accelerator (HIEPA)**

Thank you

Backup slides
