

BESIII $\pi^+\pi^-/\pi^+\pi^-\pi^0$ ISR

Yaqian WANG



JOHANNES GUTENBERG
UNIVERSITÄT MAINZ

(on behalf of the BESIII coloboration)



Johannes Gutenberg University Mainz

The 10th International Workshop on e^+e^- collisions from ϕ to ψ
23-26 Deptember, 2015
Hefei, China

Outline

- 1 Introduction
- 2 Data samples and BESIII Machine
- 3 $\pi^+\pi^-$
- 4 $\pi^+\pi^-\pi^0$
- 5 Summary

Outline

1 Introduction

2 Data samples and BESIII Machine


3 $\pi^+\pi^-$

4 $\pi^+\pi^-\pi^0$

5 Summary

g - 2


- $a_\mu = (g_\mu - 2)/2$
- $a_\mu^{\text{expe}} = 116592080 \pm 54 \pm 33 \times 10^{-11}$ at BNL
- $a_\mu^{\text{theo}} = 116591802 \pm 42 \pm 26 \times 10^{-11}$
- $a_\mu^{\text{expe}} - a_\mu^{\text{theo}} = (28.7 \pm 80) \times 10^{-11} \Rightarrow 3.6\sigma$ deviation



M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, Eur. Phys. J. C 71 1515 (2011)

Hadronic VP and muon $g - 2$

- Hadronic vacuum polarization



$$a_{\mu}^{\text{had,VP}}$$


$$a_{\mu}^{\text{SM}} = \left(\frac{g-2}{2}\right)_{\mu} = a_{\mu}^{\text{QED}} + a_{\mu}^{\text{had}} + a_{\mu}^{\text{weak}}$$


γ and leptonic

$Z, W^{\pm},$ and Higgs

$$a_{\mu}^{\text{had,LO}} = \frac{\alpha^2(0)}{3\pi^2} \int_{4m_{\pi}^2}^{\infty} ds \frac{K(s)}{s} R(s)$$

Hadronic VP and muon $g - 2$

- Hadronic vacuum polarization



- $$a_\mu^{\text{SM}} = \left(\frac{g-2}{2}\right)_\mu = a_\mu^{\text{QED}} + a_\mu^{\text{had}} + a_\mu^{\text{weak}}$$


γ and leptonic

Z, W^\pm , and Higgs

$$a_\mu^{\text{had,LO}} = \frac{\alpha^2(0)}{3\pi^2} \int_{4m_\pi^2}^{\infty} ds \frac{K(s)}{s} R(s) \rightarrow \frac{\sigma(e^+e^- \rightarrow \text{hadrons})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)}$$

Hadronic VP and muon $g - 2$

- Hadronic vacuum polarization




$$a_{\mu}^{\text{SM}} = \left(\frac{g-2}{2}\right)_{\mu} = a_{\mu}^{\text{QED}} + a_{\mu}^{\text{had}} + a_{\mu}^{\text{weak}}$$


γ and leptonic

$Z, W^{\pm},$ and Higgs

$$a_{\mu}^{\text{had,LO}} = \frac{\alpha^2(0)}{3\pi^2} \int_{4m_{\pi}^2}^{\infty} ds \frac{K(s)}{s} R(s) \longrightarrow \frac{\sigma(e^{+}e^{-} \rightarrow \text{hadrons})}{\sigma(e^{+}e^{-} \rightarrow \mu^{+}\mu^{-})}$$



ISR measurement at BaBar



D. Bernard [BaBar Collaboration], PoS Hadron **2013**, 126 (2013) [arXiv:1402.0618 [hep-ex]].


- Most important channels: $\pi^+\pi^-$, KK , $\pi^+\pi^-\pi^0$, $\pi^+\pi^-2\pi^0$
- Largest contribution to uncertainty: $\pi^+\pi^-$, $\pi^+\pi^-2\pi^0$, $KK\pi\pi$

Outline


- 1 Introduction
- 2 Data samples and BESIII Machine**
- 3 $\pi^+\pi^-$
- 4 $\pi^+\pi^-\pi^0$
- 5 Summary

BEPCII


- τ -charm factory
- Beam energy: 2 - 4.6 GeV
- Design luminosity:
 $10^{33} \text{ cm}^{-2} \text{ s}^{-1}$ (at 3.773 GeV)
- Linac + double storage ring




BESIII Detector



Data samples




Data samples



Outline

- 1 Introduction
- 2 Data samples and BESIII Machine
- 3 $\pi^+\pi^-$
- 4 $\pi^+\pi^-\pi^0$
- 5 Summary




$\pi^+\pi^-$ at BaBar and KLOE



- Obvious discrepancy between BaBar and KLOE
- High precision measurement @ **BESIII**

Event Selection and Particle Identification


- Kinematic Fit for $\pi^+\pi^-\gamma_{ISR}$
- MDC, TOF, and EMC for electron rejection
- Artificial Neuronal Network for $\mu - \pi$ separation




Systematic Uncertainties

| Source | Uncertainty (%) |
|---|-----------------|
| Photon efficiency | 0.2 |
| Tracking efficiency | 0.3 |
| Pion ANN efficiency | 0.2 |
| Pion e-PID efficiency | 0.2 |
| Angular acceptance | 0.1 |
| Background subtraction | 0.1 |
| Unfolding | 0.2 |
| FSR correction δ_{FSR} | 0.2 |
| Vacuum polarization correction δ_{vac} | 0.2 |
| Radiator function | 0.5 |
| Luminosity \mathcal{L} | 0.5 |
| Sum | 0.9 |

$\pi^+\pi^-$ Cross Section






- $\sigma^{\text{bare}}(\sqrt{s'}) = \frac{1}{\frac{2\sqrt{s'}}{s} W(s,x) \epsilon(\sqrt{s'}) \mathcal{L} \delta_{vac} \delta_{FSR}} \frac{dN}{d\sqrt{s'}}$
- ρ - ω interference clearly visible

$\pi^+\pi^-$ Form Factor (Gounaris-Sakurai Parameterization)


| Parameter | BESIII value | PDG 2014 |
|--------------------------|-----------------|-------------------|
| m_ρ [MeV/ c^2] | 774.8 ± 0.4 | 775.26 ± 0.25 |
| Γ_ρ [MeV] | 151.1 ± 0.7 | 147.8 ± 0.9 |
| m_ω [MeV/ c^2] | 782.1 ± 0.6 | 782.65 ± 0.12 |
| Γ_ω [MeV] | fixed to PDG | 8.49 ± 0.08 |
| $ c_\rho $ [10^{-3}] | 1.7 ± 0.2 | - |
| $ \phi_\omega $ [rad] | 0.04 ± 0.13 | - |

Comparison to Other $\pi^+\pi^-$ Measurements



- New BESIII measurement agrees with KLOE and BaBar
- Small shift wrt. BaBar above ρ - ω interference

Final Result: Contribution to $a_\mu^{\text{VP,LO}}$






- Precision competitive with previous measurements
- BESIII measurement between BaBar and KLOE
- $a_\mu^{\pi\pi, \text{LO}}(600 - 900 \text{ MeV}) = (370.0 \pm 2.5_{\text{stat}} \pm 3.3_{\text{sys}}) \cdot 10^{-10}$
- Confirms deviation of 3.4σ between experiment and theory
- arXiv:1507.08188 and submitted to PLB

Outline


- 1 Introduction
- 2 Data samples and BESIII Machine
- 3 $\pi^+\pi^-$
- 4 $\pi^+\pi^-\pi^0$
- 5 Summary


$$e^+e^- \rightarrow \pi^+\pi^-\pi^0$$

- History of σ for $e^+e^- \rightarrow \pi^+\pi^-\pi^0$:
 - $\sqrt{s} \lesssim 1$ GeV: $\omega(782)$ and $\phi(1020)$
 - Published results above ϕ :
 - SND : up to 1.4 GeV
 - DM2 : 1.34 ~ 2.40 GeV
 - BaBar : 1.05 ~ 3.00 GeV



 ω  ϕ  ω' and ω''



Belle and SND





$$e^+e^- \rightarrow \gamma_{\text{ISR}}\pi^+\pi^-\pi^0 \text{ from Belle}$$


$$e^+e^- \rightarrow \pi^+\pi^-\pi^0 \text{ from SND}$$

$e^+e^- \rightarrow \gamma_{\text{ISR}} \pi^+\pi^-\pi^0$ at BESIII



$e^+e^- \rightarrow \gamma_{\text{ISR}} \pi^+\pi^-\pi^0$ at BESIII

- Tagged is necessary in low mass range

$e^+e^- \rightarrow \gamma_{\text{ISR}} \pi^+\pi^-\pi^0$ at BESIII

- Tagged is necessary in low mass range
- Untagged is more efficient in high mass range

$e^+e^- \rightarrow \gamma_{\text{ISR}}\pi^+\pi^-\pi^0$ at BESIII



- Tagged is necessary in low mass range
- Untagged is more efficient in high mass range
- **Both** tagged and untagged are feasible at **BESIII**. Our goal: $< 5\%$

Outline

- 1 Introduction
- 2 Data samples and BESIII Machine
- 3 $\pi^+\pi^-$
- 4 $\pi^+\pi^-\pi^0$
- 5 Summary**

Summary

- $e^+e^- \rightarrow \pi^+\pi^-$
 - Cross section is measured at BESIII with sys. below 1%
 - Δa_μ is confirmed
- $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
 - Feasible study at BESIII
 - Benefit from both tagged and untagged
- Outlook
 - Extend tagged $\pi^+\pi^-$ ISR study to threshold region
 - Untagged ISR for $\pi^+\pi^-$ cross section at higher mass range
 - Analyze $\pi^+\pi^-$ form factor from R-scan data
(130 points, $\mathcal{L} \approx 1.3\text{fb}^{-1}$)
 - Ongoing Analysis of $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ and $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$

Thank you very much!

Theoretical calculation of a_μ

$$a_\mu^{theo} = a_\mu^{QED} + a_\mu^{weak} + a_\mu^{QCD}$$

$$a_\mu^{QED} = (116584718.104 \pm 0.148) \times 10^{-11}$$

$$a_\mu^{weak} = (153.2 \pm 1.0 \pm 1.5) \times 10^{-11}$$

$$a_\mu^{QCD} = a_\mu^{LbL} + a_\mu^{VP,LO} + a_\mu^{VP,HO}$$

$$a_\mu^{VP,LO} = (6949.1 \pm 42.7) \times 10^{-11}$$

$$a_\mu^{VP,HO} = (-97.9 \pm 0.9) \times 10^{-11}$$

$$a_\mu^{LbL} = (105 \pm 26) \times 10^{-11} \quad (\text{Glasgow consensus})$$

