

(Charmonium) Spectroscopy with BESIII

Wolfgang Gradl

on behalf of the BESIII collaboration

Athos 3/ PWA 8
Ashburn, 17th April 2015



THE LOW-ENERGY FRONTIER
OF THE STANDARD MODEL



JOHANNES GUTENBERG
UNIVERSITÄT MAINZ



PRISMA



Outline

- Introduction: charmonium spectroscopy
- New (conventional) charmonium state: $\psi_2(1^3D_2)$
- Exotic charmonium(-like) states: the X, Y, & Z
- Summary and Outlook

Not covered in this talk:

Baryon spectroscopy: see Beijiang Liu's talk on Tuesday

Light hadrons: partially covered by Andrzej Kupsc's talk

QCD bound systems

States found in nature: colour-neutral combinations

We know

mesons and baryons



QCD also allows

molecules/multi-quarks



hybrids



glueballs



and more

Totalitarian principle of quantum mechanics:

Everything not forbidden is compulsory

Multi-quark states: seen on page 1 of the quark model

Volume 8, number 3

PHYSICS LETTERS

1 February 1964



A SCHEMATIC MODEL OF BARYONS AND MESONS *

M. GELL-MANN

California Institute of Technology, Pasadena, California

Received 4 January 1964

If we assume that the strong interactions of baryons and mesons are correctly described in terms of the broken "eightfold way" ¹⁻³, we are tempted to look for some fundamental explanation of the situation. A highly promised approach is the purely dynamical "bootstrap" model for all the strongly interacting particles within which one may try to derive isotopic spin and strangeness conservation and broken eightfold symmetry from self-consistency alone ⁴. Of course, with only strong interactions, the orientation of the asymmetry in the unitary space cannot be specified; one hopes that in some way the selection of specific components of the F-spin by electromagnetism and the weak interactions determines the choice of isotopic spin and hypercharge directions.

Even if we consider the scattering amplitudes of strongly interacting particles on the mass shell only and treat the matrix elements of the weak, electromagnetic, and gravitational interactions by means of dispersion theory, there are still meaningful and important questions regarding the algebraic proper-

ties. The most interesting example of such a model is one in which the triplet has spin $\frac{1}{2}$ and $z = -1$, so that the four particles d^- , s^- , u^0 and b^0 exhibit a parallel with the leptons.

A simpler and more elegant scheme can be constructed if we allow non-integral values for the charges. We can dispense entirely with the basic baryon b if we assign to the triplet t the following properties: spin $\frac{1}{2}$, $z = -\frac{1}{3}$, and baryon number $\frac{1}{3}$. We then refer to the members $u^{\frac{2}{3}}$, $d^{-\frac{1}{3}}$, and $s^{-\frac{1}{3}}$ of the triplet as "quarks" ⁶ q and the members of the anti-triplet as anti-quarks \bar{q} . Baryons can now be constructed from quarks by using the combinations (qqq) , $(qqq\bar{q})$, etc., while mesons are made out of $(q\bar{q})$, $(qq\bar{q}\bar{q})$, etc. It is assuming that the lowest baryon configuration (qqq) gives just the representations **1**, **8**, and **10** that have been observed, while the lowest meson configuration $(q\bar{q})$ similarly gives just **1** and **8**.

A formal mathematical model based on field theory can be built up for the quarks exactly as for

Where are they?

The absence of exotics is one of the most obvious features of QCD.

R. Jaffe, hep-ph/0409065

The story of the pentaquark shows how poorly we understand QCD.

attributed to F. Wilczek, see T. Barnes, hep-ph/0510365

Where are they?


The absence of exotics is one of the most obvious features of QCD.

R. Jaffe, hep-ph/0409065


The story of the pentaquark shows how poorly we understand QCD.

attributed to F. Wilczek, see T. Barnes, hep-ph/0510365


in the past few years, compelling evidence for states beyond simple $q\bar{q}$!




$$e^+e^- \rightarrow \pi^- \pi^+ J/\psi$$




$$e^+e^- \rightarrow \pi^0 \pi^0 J/\psi$$




$$e^+e^- \rightarrow \pi^- \pi^+ h_c$$



$$e^+e^- \rightarrow \pi^0 \pi^0 h_c$$



$$e^+e^- \rightarrow \pi^- (D\bar{D}^*)^+$$




$$e^+e^- \rightarrow \pi^- (D^* \bar{D}^*)^+$$

Charmonium Spectroscopy


Charmonium and charmonium-like states useful for this search:

- $m_c \approx 1.3 \text{ GeV}$: probe transition region from perturbative to non-perturbative regime
- separation between states larger
- states presumably less mixed than in light quark sector
- can be produced copiously in e^+e^- collisions
- Exciting possibility to find exotics among new states



Godfrey & Isgur,
Phys. Rev. D **32**, 189 (1985)

Charmonium spectrum



Charmonium: $c\bar{c}$

Example potential

$$V_0^{c\bar{c}} = -\frac{4}{3} \frac{\alpha_s}{r} + br + \frac{32\pi\alpha_s}{9m_c^2} \delta(r) \vec{S}_c \vec{S}_{\bar{c}}$$

$$V_{\text{spin-dep.}} = \frac{1}{m_c^2} \left[\left(\frac{2\alpha_s}{r^3} - \frac{b}{2r} \right) \vec{L} \cdot \vec{S} + \frac{4\alpha_s}{r^3} T \right]$$


+ relativistic corrections!

Godfrey & Isgur, PRD 32, 189 (1985);
Barnes, Godfrey & Swanson,
PRD 72, 054026 (2005)

Use well-established states to fix parameters, then predict remainder of spectrum, and transitions

➔ Remarkably good description above $D\bar{D}$ threshold: some mass shifts

Charmonium spectrum



Charmonium: $c\bar{c}$

Example potential

$$V_0^{c\bar{c}} = -\frac{4}{3} \frac{\alpha_s}{r} + br + \frac{32\pi\alpha_s}{9m_c^2} \delta(r) \vec{S}_c \vec{S}_{\bar{c}}$$

$$V_{\text{spin-dep.}} = \frac{1}{m_c^2} \left[\left(\frac{2\alpha_s}{r^3} - \frac{b}{2r} \right) \vec{L} \cdot \vec{S} + \frac{4\alpha_s}{r^3} T \right]$$

+ relativistic corrections!

Godfrey & Isgur, PRD 32, 189 (1985);
Barnes, Godfrey & Swanson,
PRD 72, 054026 (2005)

Use well-established states to fix parameters, then predict remainder of spectrum, and transitions

➔ Remarkably good description above $D\bar{D}$ threshold: some mass shifts



BESIII: a τ -charm factory

BEPCII and BESIII




Linac

BESIII


BSRF

Tiananmen 10km

A τ -charm factory






BESIII detector



Completely new detector

Comparable performance to CLEO-c, + muon ID

BESIII data sets




+ 104 energy points between 3.85 and 4.59 GeV
 + ~ 20 energy points between 2.0 and 3.1 GeV
 (*ongoing*)

Direct production of 1^{--} states studied
 with world's largest scan dataset




A new conventional $c\bar{c}$ state

Higher charmonium states



The X(3823) at Belle

PRL **111**, 032001 (2013)



Using full Belle data set of
 $772 \times 10^6 B\bar{B}$

$B \rightarrow K\gamma\chi_{c1}$
simultaneous fit to B^+ and B^0

3.8σ evidence

$M = 3823.1 \pm 1.8 \pm 0.7 \text{ MeV}$
very narrow


Mass (and width) compatible with
 $\psi_2(1^3D_2)$ state

$$e^+e^- \rightarrow \pi^+\pi^-\chi(3823) \rightarrow \pi^+\pi^-\gamma\chi_{c1}$$

reconstruct $\chi_{c1,2} \rightarrow \gamma J/\psi \rightarrow \gamma \ell^+ \ell^-$

look in mass recoiling against $\pi^+\pi^-$ system, $M_{\text{recoil}}(\pi^+\pi^-)$

Use 5 large data sets (total luminosity $\sim 4.1 \text{ fb}^{-1}$)







$$e^+e^- \rightarrow \pi^+\pi^-\chi(3823) \rightarrow \pi^+\pi^-\gamma\chi_{c1}$$

$$\sqrt{s} \geq 4.36 \text{ GeV}$$

$$\sqrt{s} = 4.23, 4.26 \text{ GeV preliminary}$$

arXiv:1503.08203




$$M = 3821.7 \pm 1.3 \pm 0.7 \text{ MeV, significance } 6.7\sigma$$

$$\Gamma < 16 \text{ MeV at } 90\% \text{ C.L.}$$

$$e^+e^- \rightarrow \pi^+\pi^-X(3823) \rightarrow \pi^+\pi^-\gamma\chi_{c1}$$

Energy-dependent cross section for

$$e^+e^- \rightarrow \pi^+\pi^-X(3823) \rightarrow \pi^+\pi^-\gamma\chi_{c1}$$



Mass and width \sim in agreement with potential model prediction for 1^3D_2 predicted to be narrow!

Production ratio


$$R_{21} \equiv \frac{\mathcal{B}(X(3823) \rightarrow \gamma\chi_{c2})}{\mathcal{B}(X(3823) \rightarrow \gamma\chi_{c1})}$$

~ 0.2 prediction
 < 0.43 at 90% C.L.

Compatible with both $Y(4360)$ and $\psi(4415)$ line shapes

$$e^+e^- \rightarrow \pi^+\pi^-\chi(3823) \rightarrow \pi^+\pi^-\gamma\chi_{c1}$$

Angular distribution $\theta \equiv \angle(\pi\pi, \psi_2)$
 assuming $\pi\pi$ system in S -wave: $1 + \cos^2\theta$ for spin 2




Not enough statistics to distinguish S and D wave
 from data

Mass and width \sim in agreement
 with potential model prediction for
 1^3D_2
 predicted to be narrow!

J^P by exclusion:
 $1^1D_2 \rightarrow \gamma\chi_{c1}$ forbidden
 $1^3D_3 \rightarrow \gamma\chi_{c1}$ has zero amplitude

Good candidate for $\psi_2(1^3D_2)$


Higher charmonium states — a new family member!






Exotic states: the X and Y

Surprising discoveries: the XYZ states




Most of the 'XYZ' states discovered at Belle and BABAR in e^+e^- collisions in bottomonium region e.g. in B decays:




Surprising discoveries: the XYZ states

Most of the 'XYZ' states discovered at Belle and BABAR in e^+e^- collisions in bottomonium region




$$B^+ \rightarrow K^+ \pi^+ \pi^- J/\psi$$




Belle, PRL 91, 262001 (2003)

The X(3872)




Extremely narrow, sits at or just below the $D\bar{D}^*$ threshold



$$M = 3871.69 \pm 0.17 \text{ MeV}/c^2$$

$$\Gamma < 1.2 \text{ MeV}$$

The $Y(4260) \rightarrow J/\psi \pi^+ \pi^-$




e^+e^- collisions near $Y(4S)$


in ISR production

$$e^+e^- \rightarrow \gamma_{\text{ISR}} J/\psi \pi^+ \pi^-$$

$$\Rightarrow J^{PC} = 1^{--}$$



The $Y(4260) \rightarrow J/\psi \pi^+ \pi^-$




e^+e^- collisions near $Y(4S)$

in ISR production


$e^+e^- \rightarrow \gamma_{\text{ISR}} J/\psi \pi^+ \pi^-$

$\Rightarrow J^{PC} = 1^{--}$



BABAR, PRD 86, 051102(R) (2012)

The $Y(4260) \rightarrow J/\psi \pi^+ \pi^-$




... $Y(4260) \rightarrow J/\psi \pi^+ \pi^-$

... $Y(4360) \rightarrow \psi(2S) \pi^+ \pi^-$

... additional state at 4660 MeV





■ supernumerary states:
all 1^{--} slots already taken

➔ do not correspond to peaks in
 $\sigma(e^+e^- \rightarrow \text{hadrons})$





➔ produce them directly at BESIII!

$$e^+e^- \rightarrow \gamma X(3872) \rightarrow \gamma J/\psi \pi^+ \pi^-$$

BESIII, PRL **112**, 092001 (2014)

Clear ISR ψ' signal for validation
 $X(3872)$ signal around 4.23 – 4.26 GeV

$$e^+e^- \rightarrow \gamma X(3872) \rightarrow \gamma J/\psi \pi^+ \pi^-$$



 BESIII, PRL **112**, 092001 (2014)

 20.1 ± 4.5 events

 significance 6.3σ
 $M = 3871.9 \pm 0.7 \pm 0.2 \text{ MeV}/c^2$

[PDG2013:

 $3871.68 \pm 0.17 \text{ MeV}/c^2$

$$e^+e^- \rightarrow \gamma X(3872) \rightarrow \gamma J/\psi \pi^+ \pi^-$$



Suggestive of radiative transition
Y(4260) \rightarrow γ X(3872)




Direct connection between the two
states?



The Z_c family

$e^+e^- \rightarrow J/\psi \pi^+ \pi^-$ at 4.26 GeV

BESIII, PRL **110**, 252001 (2013)






...have hundreds of events!



$J/\psi \pi^+ \pi^-$ Dalitz plot






BESIII, PRL **110**, 252001 (2013)



Model $\pi^+\pi^-$ -system with known structure:
 $f_0(500)$, $f_0(980)$, non-resonant
obtain good fit of $\pi^+\pi^-$ mass projection



$J/\psi \pi^+ \pi^-$ Dalitz plot

BESIII, PRL **110**, 252001 (2013)



$e^+e^- \rightarrow J/\psi \pi^+ \pi^-$ at 4.26 GeV

BESIII, PRL **110**, 252001 (2013)



Charged charmonium-like structure

$$M = (3899.0 \pm 3.6 \pm 4.9) \text{ MeV}/c^2$$

$$\Gamma = (46 \pm 10 \pm 20) \text{ MeV}$$

Confirmed by Belle PRL **110**, 252002
and with CLEOC data PLB **727**, 366

Close to DD^* threshold
Interpretation?

A neutral partner to the $Z_c(3900)^+$?


If interpretation of $Z_c(3900)^+$ as four-quark state is correct:
expect state completing isospin triplet, with decay $Z_c(3900)^0 \rightarrow \pi^0 J/\psi$

A neutral partner to the $Z_c(3900)^+$?

If interpretation of $Z_c(3900)^+$ as four-quark state is correct:

expect state completing isospin triplet, with decay $Z_c(3900)^0 \rightarrow \pi^0 J/\psi$

Study $e^+e^- \rightarrow \pi^0 \pi^0 J/\psi$ with large data sets at three different \sqrt{s}




Structure in $\pi^0 J/\psi$ invariant mass clearly visible at all energies

$$M = 3894.8 \pm 2.3 \pm 2.7 \text{ MeV}/c^2$$


$$\Gamma = 29.6 \pm 8.2 \pm 8.2 \text{ MeV}$$

Significance 10σ



$Z_c(3900)^+$ at $D\bar{D}^*$ threshold

Decay mode $Z_c(3900)^+ \rightarrow (D\bar{D}^*)^+$?





$Z_c(3900)^+$ at $D\bar{D}^*$ threshold

BESIII, PRL **112**, 022001 (2014)

Decay mode $Z_c(3900)^+ \rightarrow (D\bar{D}^*)^+$?

Single tag analysis:

- reconstruct 'bachelor' π^+ and $D^0 \rightarrow K^-\pi^+$ or $D^- \rightarrow K^+\pi^-\pi^-$
- require D^* in missing mass
- veto $e^+e^- \rightarrow (D^*\bar{D}^*)^0$
- apply kinematic fit; look in mass recoiling against π^+





$Z_c(3900)^+$ at $D\bar{D}^*$ threshold

BESIII, PRL **112**, 022001 (2014)

Decay mode $Z_c(3900)^+ \rightarrow (D\bar{D}^*)^+$?


$e^+e^- \rightarrow \pi^+ D^0 D^{*-}$ at BESIII



...and BESIII sees structure in $(DD^*)^\pm$


Large systematics due to non- Z_c signal shape

$Z_c(3900)^+$ at $D\bar{D}^*$ threshold




New: Double tag analysis

- reconstruct 'bachelor' π^+ and D^0, D^- in 4 or 6 decay modes
- kinematic fit, requiring π from D^* in missing mass essentially background-free D^*
- improved statistics, much better control over background shape, improved systematics
- $M^{\text{recoil}}(\pi^+) = M(D\bar{D}^*)$



$e^+e^- \rightarrow \pi^+(D\bar{D}^*)^-$ with double tags



Simultaneous fit with phase space shape + $(BW \otimes \mathcal{R}) \times \epsilon$

Compatible with, but significantly more precise, than single-tag analysis

$$M = 3884.3 \pm 1.2 \pm 1.5 \text{ MeV}/c^2$$

$$\Gamma = 23.8 \pm 2.1 \pm 2.6 \text{ MeV}$$

$e^+e^- \rightarrow \pi^+(D\bar{D}^*)^-$ with double tags: Results

	BESIII single D tags PRL 112, 022001	BESIII double D tags preliminary
$M_{\text{pole}}[\text{MeV}/c^2]$	$3883.9 \pm 1.5(\text{stat}) \pm 4.2(\text{syst})$	$3884.3 \pm 1.2(\text{stat}) \pm 1.5(\text{syst})$
$\Gamma_{\text{pole}}[\text{MeV}]$	$24.8 \pm 3.3(\text{stat}) \pm 11.0(\text{syst})$	$23.8 \pm 2.1(\text{stat}) \pm 2.6(\text{syst})$
$\sigma \times \mathcal{B}[\text{pb}]$		
4.23 GeV		$106.8 \pm 7.1(\text{stat}) \pm 9.5(\text{syst})$
4.26 GeV	$83.5 \pm 6.6(\text{stat}) \pm 22.0(\text{syst})$	$88.0 \pm 6.1(\text{stat}) \pm 7.9(\text{syst})$

$$\sigma \times \mathcal{B} \equiv \sigma(e^+e^- \rightarrow \pi^\pm Z_c(3885)^\mp) \times \mathcal{B}(Z_c(3885)^\mp \rightarrow (D\bar{D}^*)^\mp)$$

$Z_c(3885)^+$ Quantum numbers?

θ_π : angle between bachelor pion and beam axis in CMS

Know initial state is 1^- , with $J_z = \pm 1$. Depending on J^P of Z_c :


0^+ excluded by parity conservation

0^- π and $Z_c(3885)$ in P -wave, with $J_z = \pm 1$ $\Rightarrow dN/d \cos \theta_\pi \propto 1 - \cos^2 \theta_\pi$

1^- π and $Z_c(3885)$ in P -wave $\Rightarrow dN/d \cos \theta_\pi \propto 1 + \cos^2 \theta_\pi$

1^+ π and $Z_c(3885)$ in S or D wave.

Assume D wave small near threshold: $\Rightarrow dN/d \cos \theta_\pi \propto 1$



Efficiency corrected event yield
in 10 bins in $|\cos \theta_\pi|$

data clearly favour $J^P = 1^+$
for $D\bar{D}^*$ structure

confirms J^P for $Z_c(3885)$ from single-tags



$$e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$$

Exclusively reconstruct the process

$$e^+e^- \rightarrow \pi^+\pi^-h_c(1P)$$

$$h_c(1P) \rightarrow \gamma\eta_c(1S)$$

$$\eta_c(1S) \rightarrow 16 \text{ decay channels}$$





$$e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$$

Exclusively reconstruct the process



$$e^+e^- \rightarrow \pi^+\pi^-h_c(1P)$$

$$h_c(1P) \rightarrow \gamma\eta_c(1S)$$

$$\eta_c(1S) \rightarrow 16 \text{ decay channels}$$



$$e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$$



Charged charmonium-like structure
close to $D^*\bar{D}^*$ threshold

$$M = 4022.9 \pm 0.8 \pm 2.7 \text{ MeV}/c^2$$

$$\Gamma = 7.9 \pm 2.7 \pm 2.6 \text{ MeV}$$



Note: no significant signal for
 $Z_c(3900)^+ \rightarrow \pi^+ h_c$ seen!

$$e^+e^- \rightarrow h_c(1P)\pi^0\pi^0$$

Study $e^+e^- \rightarrow \pi^0\pi^0 h_c$ at 4.23, 4.26, 4.36 GeV

Observe structure in $h_c\pi^0$ mass distribution:

Neutral partner to $Z_c(4020)^+$




$$M = 4023.6 \pm 4.5 \text{ MeV/c}^2$$

Γ fixed in the fit

Isospin triplet found!

Yet another mass threshold ...

$Z_c(4020)$ at D^*D^* threshold





Yet another mass threshold ...

BESIII, PRL **112**, 132001 (2014)

$Z_c(4020)$ at D^*D^* threshold

$e^+e^- \rightarrow \pi^+(D^*\bar{D}^*)^-$ at BESIII




...and BESIII sees structure in D^*D^*


$$M = 4026.3 \pm 2.6 \pm 3.7 \text{ MeV}/c^2$$

$$\Gamma = 24.8 \pm 5.6 \pm 7.7 \text{ MeV}$$


All the Z_c s from BESIII near $\sqrt{s} = 4.3$ GeV




$$e^+e^- \rightarrow \pi^- \pi^+ J/\psi$$




$$e^+e^- \rightarrow \pi^0 \pi^0 J/\psi$$



$$e^+e^- \rightarrow \pi^- \pi^+ h_c$$



$$e^+e^- \rightarrow \pi^0 \pi^0 h_c$$




$$e^+e^- \rightarrow \pi^- (D\bar{D}^*)^+$$

$$Z_c(3900)^+?$$

BESIII

in progress

$$Z_c(3900)^0?$$



$$e^+e^- \rightarrow \pi^- (D^* \bar{D}^*)^+$$

$$Z_c(4020)^+?$$

BESIII

in progress

$$Z_c(4020)^0?$$


Nature of these states? Isospin triplets?

Different decay channels of the same states observed?

Other decay modes?

Other decay modes?

Exploring new decay modes crucial to identify nature of structures close to threshold





Decay modes with $c\bar{c}$ annihilation does not involve hidden or open charm final states!

If $c\bar{c}$ in S -wave, annihilation should be 'easy' ...

$$Z_C(3900)^+ \rightarrow \omega\pi^+ \rightarrow (\pi^+\pi^-\pi^0)\pi^+$$

$$\sqrt{s} = 4.230 \text{ GeV}$$

$$\sqrt{s} = 4.260 \text{ GeV}$$






$$\sigma(e^+e^- \rightarrow Z_C^+ \pi^-, Z_C^+ \rightarrow \omega\pi^+) < 0.27 \text{ pb}$$

$$\sigma(e^+e^- \rightarrow Z_C^+ \pi^-, Z_C^+ \rightarrow \omega\pi^+) < 0.18 \text{ pb}$$

Compared to sum of $Z_C^+ \rightarrow J/\psi \pi^+$ and $Z_C^+ \rightarrow (D\bar{D}^*)^+$:

$$\Gamma(Z_C^+ \rightarrow \omega\pi^+) < 0.2\% \Gamma_{\text{tot}}$$

$$e^+e^- \rightarrow \eta J/\psi$$



Compare to $e^+e^- \rightarrow \gamma_{ISR} \eta J/\psi$ from Belle, Phys. Rev. D 87, 051101(R) (2013)


Good agreement, significantly better precision

Cross section peaks around 4.2 GeV

Also searched for $e^+e^- \rightarrow \pi^0 J/\psi$: no significant signal found

$$e^+e^- \rightarrow \eta J/\psi \text{ vs } e^+e^- \rightarrow \pi^+ \pi^- J/\psi$$

BESIII preliminary,
arXiv:1503.06644 [hep-ex]



Compare to $e^+e^- \rightarrow \gamma_{\text{ISR}} \pi^+ \pi^- J/\psi$ from
Belle, Phys. Rev. Lett. **110**, 252002 (2013)

Very different line shape

➡ Different dynamics at work in
 $e^+e^- \rightarrow \eta J/\psi$ compared to
 $e^+e^- \rightarrow \pi^+ \pi^- J/\psi$


Search for $Y(4140) \rightarrow J/\psi \phi$

CDF first reported evidence for
 $Y(4140) \rightarrow J/\psi \phi$ in $B^+ \rightarrow J/\psi \phi K^+$,
also claimed by D0 and CMS

Not seen by LHCb, Belle (B decays and $\gamma\gamma$ events),
or BABAR

$J/\psi \phi$ system has $C = +1$: search in radiative transitions of charmonium or $Y(4260)$

If both $Y(4260)$ and $Y(4140)$ are *charmonium hybrids*:
partial width of $Y(4260) \rightarrow \gamma Y(4140)$ may be up to several tens of keV
N. Mahajan, PLB **679**, 228 (2009)



CDF, PRL **102**, 242002, (2009)


Search for $Y(4140) \rightarrow J/\psi \phi$

Use BESIII's large data samples from 4.23 – 4.36 GeV (2.47 fb^{-1} in total)

$$e^+e^- \rightarrow \gamma J/\psi \phi$$

$$J/\psi \rightarrow e^+e^-, \mu^+\mu^-,$$

$$\phi \rightarrow K^+K^-, K_S^0K_L^0, \pi^+\pi^-\pi^0$$



Search for $Y(4140) \rightarrow J/\psi \phi$

No significant signal found; place upper limits on
 $\sigma(e^+e^- \rightarrow \gamma Y(4140)) \times \mathcal{B}(Y(4140) \rightarrow J/\psi \phi)$


Compare sensitivity to $e^+e^- \rightarrow \gamma X(3872) \times \mathcal{B}(X(3872) \rightarrow J/\psi \pi^+ \pi^-)$

\sqrt{s}/GeV	4.23	4.26	4.36
$\sigma \times \mathcal{B}(X(3872))/\text{pb}$	0.27 ± 0.09	0.33 ± 0.12	0.11 ± 0.09
$\sigma \times \mathcal{B}(Y(4140))/\text{pb}$	< 0.35	< 0.28	< 0.33

Assuming $\mathcal{B}(Y(4140) \rightarrow J/\psi \phi) \sim 30\%$ and $\mathcal{B}(X(3872) \rightarrow J/\psi \pi^+ \pi^-) \sim 5\%$:


$$\frac{\sigma[e^+e^- \rightarrow \gamma Y(4140)]}{\sigma[e^+e^- \rightarrow \gamma X(3872)]} < 0.1 \quad \text{at 4.23, 4.26 GeV}$$

Summary



- Quark model describes charmonium states $c\bar{c}$ reasonably well
- XYZ states: unexpected, point to non-conventional states ($c\bar{c}g$, $cq\bar{q}c$, $(\bar{c}q)(\bar{q}c)$, $c\bar{c}\pi\pi$...)
- Observation of transitions between XYZ states
- ➔ Start making connections between new, exotic states
- ➔ Dynamically generated at thresholds, or new kind of QCD bound states?

Summary




- Structure of XYZ to be clarified; learn more about strongly bound systems
- More detailed studies (PWA, other channels ...) at BESIII ongoing
- Future:
More data from BESIII
LHCb spectroscopy
Belle-II will start 2017
- Exciting times ahead



謝
謝

!

BEPCII storage rings: a τ -charm factory



Upgrade of BEPC (started 2004,
first collisions July 2008)

Beam energy **1 ... 2.3 GeV**

Optimum energy **1.89 GeV**

Single beam current **0.91 A**

Crossing angle **± 11 mrad**

Design luminosity **$10^{33} \text{ cm}^{-2} \text{ s}^{-1}$**

Achieved **$8 \times 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$**

Beam energy measurement:

Laser Compton backscattering


$\Delta E/E \approx 5 \times 10^{-5}$

(≈ 50 keV at τ threshold)

Kinematic reflections

In multi-body decays, resonance in one subchannel can produce peaks in other mass projections (**reflections**)

For example $D^0 \rightarrow K^+ K^- \pi^0$: relatively easy to understand







Kinematic reflections

But can be much less obvious



Example: high-statistics analysis of decay $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ BABAR, PRD78,034023 (2008)

$$\mathcal{A}_D(m_-^2, m_+^2) = \sum_r a_r e^{i\phi_r} \mathcal{A}_r(m_-^2, m_+^2) + a_{\text{NR}} e^{i\phi_{\text{NR}}}$$

Using 10 resonant amplitudes



The B factories Belle and BABAR




mainly $e^+e^- \rightarrow Y(4S) \rightarrow B\bar{B}$
Asymmetric beam energies


- KEK-B: $8 \text{ GeV } e^- \times 3.5 \text{ GeV } e^+$
- $\mathcal{L}_{\text{int}} \approx 1 \text{ ab}^{-1}$
- Data taking finished 2010

- PEP-II: $9 \text{ GeV } e^- \times 3.1 \text{ GeV } e^+$
- $\mathcal{L}_{\text{int}} \approx 530 \text{ fb}^{-1}$
- Data taking finished 2008

Spin-Parity assignment



Phase motion



behaves like a 'true' resonance

$$J^{PC} = 1^{++} \text{ preferred}$$