

Charm Physics at BESIII

Peilian LIU

On behalf of the BESIII Collaboration

Institute of High Energy Physics, CAS


liupl@ihep.ac.cn



Outline

- **BEPCII Collider and BESIII Detector**
- **Data samples of charmed hadrons**
- **Λ_c^+ decays**
 - Λ_c^+ hadronic decays
 - Λ_c^+ semi-leptonic decays
- **D decays**
 - D hadronic decays
 - D leptonic and semi-leptonic decays
- **Summary**

Beijing Electron Positron Collider (BEPCII)




- ✓ Beam energy: 1.0 – 2.3 GeV
- ✓ Luminosity reached the design value (04/05/2016)

$$1.0 \times 10^{33} \text{cm}^{-2} \text{s}^{-1}$$

1.0 Tesla super-conducting magnet

BESIII Detector



Be beam pipe

Muon Counters

9/8 RPC layers (barrel/endcaps)

Cut-off momentum: 0.4 GeV/c

Csl(Tl) ElectroMagnetic Calorimeter

σ_E/E (at 1 GeV): 2.3%

$\sigma_{z,\phi}$ (at 1 GeV) 5 ~ 7 mm

Time of Flight (TOF)


σ_T : 68/100 ps (barrel/endcaps)

Drift Chamber(MDC)

σ_p/p (at 1 GeV): 0.32%

$\sigma_{dE/dx} < 5\%$ (Bhabha)

Data samples of charmed meson and baryon




- ✓ 2.93 fb⁻¹ data@3.773 GeV for $D^0 \bar{D}^0 / D^+ D^-$ production
- ✓ 0.48 fb⁻¹ data@4.009 GeV for $D_s^+ D_s^-$ production
- ✓ 0.57 fb⁻¹ data@4.599 GeV for $\Lambda_c^+ \bar{\Lambda}_c^-$ production

Charm physics at BESIII: Motivation

- Unitarity test of CKM matrix: measuring $|V_{cs}|$ and $|V_{cd}|$
- Lattice QCD calibration: $f_{D \rightarrow K/\pi}(q^2)$ and other formfactors, $f_{D(s)^+}$ decay constant
- New Physics: finding evidence of CP violation, rare decays, significant deviations from CKM unitarity or from LQCD calculations, $D - \bar{D}$ mixing
- Providing inputs for b-physics

Measurement of Absolute Branching Fractions

Illustration: $e^+e^- \rightarrow \psi(3770) \rightarrow D\bar{D}$




- **Single Tag (ST)**
 - ✓ Tag the charmed meson or baryon via hadronic decays with large Branching Fractions (BF)
 - ✓ $\Delta E \equiv E_{rec.} - E_{beam}$
 - ✓ $M_{BC}^2 c^4 \equiv E_{beam}^2 - p^2 c^2$
- **Double Tag (DT)**
 - ✓ Reconstruct signals in the recoil side against
 - ✓ For (semi)leptonic decays: $U_{miss} \equiv E_{miss} - p_{miss}$
- **Absolute BF** $BF(D \rightarrow sig) = \frac{N_{sig}}{N_{tag} \times \epsilon_{sig} / \epsilon_{tag, sig}}$.
only need the yields (N) and the efficiencies(ϵ) of ST and DT

The advantage of data at threshold

- ✓ $D_{(s)}\bar{D}_{(s)}/\Lambda_c^+\bar{\Lambda}_c^-$ pairs produced at threshold, no additional hadrons
- ✓ Effectively suppress the background with the DT technique
- ✓ Enable the measurement of absolute BF, without knowing the number of $D\bar{D}$ pairs
- ✓ Most systematic uncertainty in tag side are cancelled out.

Absolute hadronic BFs of Λ_c^+ baryon

□ ST events




ST yield of $\bar{\Lambda}_c^- \rightarrow \alpha$


$$N_{ST}^\alpha = N_{\Lambda_c^+ \bar{\Lambda}_c^-} \cdot BF_\alpha \cdot \epsilon_{ST}^\alpha$$

□ Extraction of the 12 BFs: Simultaneously did 24 fits

- ✓ BFs are constraint to common variable $N_{\Lambda_c^+ \bar{\Lambda}_c^-}$
- ✓ Considering statistical and systematic correlations among the different hadronic modes
- ✓ $\chi^2/\text{ndf} = 9.9/(24-13) = 0.9$



□ DT events



Total DT yield of $\Lambda_c^+ \rightarrow s$ over 12 $\bar{\Lambda}_c^- \rightarrow \alpha$

$$N_{DT}^s = N_{\Lambda_c^+ \bar{\Lambda}_c^-} \cdot BF_s \sum_{\alpha} BF_\alpha \cdot \epsilon_{DT}^\alpha$$

Absolute hadronic BFs of Λ_c^+ baryon

- The first absolute measurement of the Λ_c^+ BFs at the $\bar{\Lambda}_c^+ \bar{\Lambda}_c^-$ production threshold, since the Λ_c^+ discovery 30 years ago **PRL116, 052001 (2016)**



Mode	This work (%)	PDG (%)	BELLE \mathcal{B}
pK_S^0	$1.52 \pm 0.08 \pm 0.03$	1.15 ± 0.30	
$pK^- \pi^+$	$5.84 \pm 0.27 \pm 0.23$	5.0 ± 1.3	$6.84 \pm 0.24^{+0.21}_{-0.27}$
$pK_S^0 \pi^0$	$1.87 \pm 0.13 \pm 0.05$	1.65 ± 0.50	
$pK_S^0 \pi^+ \pi^-$	$1.53 \pm 0.11 \pm 0.09$	1.30 ± 0.35	
$pK^- \pi^+ \pi^0$	$4.53 \pm 0.23 \pm 0.30$	3.4 ± 1.0	
$\Lambda \pi^+$	$1.24 \pm 0.07 \pm 0.03$	1.07 ± 0.28	
$\Lambda \pi^+ \pi^0$	$7.01 \pm 0.37 \pm 0.19$	3.6 ± 1.3	
$\Lambda \pi^+ \pi^- \pi^+$	$3.81 \pm 0.24 \pm 0.18$	2.6 ± 0.7	
$\Sigma^0 \pi^+$	$1.27 \pm 0.08 \pm 0.03$	1.05 ± 0.28	
$\Sigma^+ \pi^0$	$1.18 \pm 0.10 \pm 0.03$	1.00 ± 0.34	
$\Sigma^+ \pi^+ \pi^-$	$4.25 \pm 0.24 \pm 0.20$	3.6 ± 1.0	
$\Sigma^+ \omega$	$1.56 \pm 0.20 \pm 0.07$	2.7 ± 1.0	

- ✓ The precisions are improved by factors of 3~6
- ✓ The golden mode $\Lambda_c^+ \rightarrow pK^- \pi^+$
 - Our measurement is consistent with the PDG value, but lower than Belle's with 2σ significance
 - Improved absolute BF of $pK^- \pi^+$ together with Belle's result are key to calibrate other decays

Absolute BF of $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$

PRL115, 221805 (2015)


11 ST modes except $\Lambda_c^+ \rightarrow \Sigma^+ \omega$



$$\text{BF}(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e) = (3.63 \pm 0.38 \pm 0.20)\%$$

- ◆ The first absolute measurement
- ◆ Improvement of the precision of the PDG value ($2.9 \pm 0.5\%$)
- ◆ Important for the testing and calibration of the LQCD calculations

Observation of $D^+ \rightarrow \omega\pi^+$ and Evidence for $D^0 \rightarrow \omega\pi^0$



Charged D tag modes:

- (a) $K^+\pi^-\pi^-$
- (b) $K^+\pi^-\pi^-\pi^0$
- (c) $K_S^0\pi^-$
- (d) $K_S^0\pi^-\pi^0$
- (e) $K_S^0\pi^+\pi^-\pi^-$
- (f) $K^+K^-\pi^-$


Neutral D tag modes:

- (g) $K^+\pi^-$
- (h) $K^+\pi^-\pi^0$
- (i) $K^+\pi^-\pi^+\pi^-$
- (j) $K^+\pi^-\pi^0\pi^0$
- (k) $K^+\pi^-\pi^+\pi^-\pi^0$

Observation of $D^+ \rightarrow \omega\pi^+$ and Evidence for $D^0 \rightarrow \omega\pi^0$

Suppress background via DT method

PRL116, 082001 (2016)




- Predictions of $\text{BF}(D \rightarrow \omega\pi) \sim 1.0 \times 10^{-4}$
- Studied by CLEO-c with ST method \rightarrow upper limit

Mode	This work	Previous measurements
$D^+ \rightarrow \omega\pi^+$	$(2.79 \pm 0.57 \pm 0.16) \times 10^{-4}$	$< 3.4 \times 10^{-4}$ at 90% C.L.
$D^0 \rightarrow \omega\pi^0$	$(1.17 \pm 0.34 \pm 0.07) \times 10^{-4}$	$< 2.6 \times 10^{-4}$ at 90% C.L.
$D^+ \rightarrow \eta\pi^+$	$(3.07 \pm 0.22 \pm 0.13) \times 10^{-3}$	$(3.53 \pm 0.21) \times 10^{-3}$
$D^0 \rightarrow \eta\pi^0$	$(0.65 \pm 0.09 \pm 0.04) \times 10^{-3}$	$(0.68 \pm 0.07) \times 10^{-3}$

- ✓ Improved understanding of U-spin and SU(3)-flavor symmetry breaking effects in D decays

Study of $D^+ \rightarrow \omega e^+ \nu_e$ and Search for $D^+ \rightarrow \phi e^+ \nu_e$

PRD92, 071101R(2015)



Dots: data


Arrows: signal region

Mode	This work	Previous
$\omega e^+ \nu_e$	$(1.63 \pm 0.11 \pm 0.08) \times 10^{-3}$	$(1.82 \pm 0.18 \pm 0.07) \times 10^{-3}$
$\phi e^+ \nu_e$	$< 1.3 \times 10^{-5}$ (90%C.L.)	$< 9.0 \times 10^{-5}$ (90%C.L.)

- No significant excess of $D^+ \rightarrow \phi e^+ \nu_e$ is observed
- More precise BFs

Form factors measurement of $D^+ \rightarrow \omega e^+ \nu_e$


PRD92, 071101R(2015)



$$\frac{d\Gamma}{dq^2 d \cos \theta_1 d \cos \theta_2 d \chi dm_{\pi\pi\pi}} = \mathcal{F}(V(q^2), A_{1,2}(q^2) \dots)$$

A five-dimensional maximum likelihood fit is performed in the space of m^2 , q^2 , $\cos\theta_1$, $\cos\theta_2$ and χ

■ Amplitude analysis of $D^+ \rightarrow \omega e^+ \nu$ is performed for the first time




$$r_V = V(0)/A_1(0) = 1.24 \pm 0.09 \pm 0.06$$

$$r_2 = A_2(0)/A_1(0) = 1.06 \pm 0.15 \pm 0.05$$



Study of dynamics of $D^0 \rightarrow (K^-/\pi^-)e^+\nu_e$

PRD92,072012(2015)



$$\text{BF}(D^0 \rightarrow K^- e^+ \nu_e) = (3.505 \pm 0.014 \pm 0.033)\%$$

$$\text{BF}(D^0 \rightarrow \pi^- e^+ \nu_e) = (0.295 \pm 0.004 \pm 0.003)\%$$



$$\frac{d\Gamma}{dq^2} = \frac{G_F^2}{24\pi^3} |V_{cs(d)}|^2 |\vec{p}_{K^-(\pi^-)}|^3 |f_+^{K(\pi)}(q^2)|^2$$

Simple pole model:

$$f_+(q^2) = \frac{f_+(0)}{1 - \frac{q^2}{M_{\text{pole}}^2}}$$

Modified pole model:

$$f_+(q^2) = \frac{f_+(0)}{(1 - \frac{q^2}{M_{\text{pole}}^2})(1 - \alpha \frac{q^2}{M_{\text{pole}}^2})}$$

Series expansion:



$$f_+(q^2) = \frac{1}{P(t)\Phi(t, t_0)} a_0(t_0) \times (1 + r_1(t_0)[z(t, t_0)])$$

Study of dynamics of $D^0 \rightarrow (K^-/\pi^-)e^+\nu_e$

PRD92,072012(2015)

$$|V_{cs(d)}| f_+^{D^0 \rightarrow (K^-/\pi^-)e^+\nu_e}(0) \rightarrow f_+^{D^0 \rightarrow (K^-/\pi^-)e^+\nu_e}(0)$$

Input $V_{cs(d)}$ of CKM Fitter




The most precise form factor measurement to calibrate the LQCD

Decay dynamics and CP asymmetry in $D^+ \rightarrow K_L^0 e^+ \nu_e$

- Regardless of long flight distance, K_L^0 interact with EMC and deposit part of energy, thus giving position information
- After reconstructing all other particles, K_L^0 can be inferred with position information and constraint $U_{\text{miss}} \rightarrow 0$

PRD92, 112008(2015)

■ Simultaneous fit to observed numbers of DT candidates



$$\text{BF}(D^+ \rightarrow K_L e^+ \nu_e) = (4.454 \pm 0.038 \pm 0.102)\%$$

$$\text{BF}(D^- \rightarrow K_L e^- \bar{\nu}_e) = (4.507 \pm 0.038 \pm 0.104)\%$$

$$|V_{cs}| = 0.975 \pm 0.008 \pm 0.015 \pm 0.025 \text{ (with LQCD input } f_+^K(0)\text{), consistent with } 0.986 \pm 0.016 \text{ in PDG}$$



✓ The first measurement of $\text{BF}(D^+ \rightarrow K_L e^+ \nu_e)$

$$A_{CP} \equiv \frac{\text{BF}(D^+ \rightarrow K_L e^+ \nu_e) - \text{BF}(D^- \rightarrow K_L e^- \bar{\nu}_e)}{\text{BF}(D^+ \rightarrow K_L e^+ \nu_e) + \text{BF}(D^- \rightarrow K_L e^- \bar{\nu}_e)} = (-0.59 \pm 0.60 \pm 1.48)\%$$

Study of $D^+ \rightarrow K^- \pi^+ e^+ \nu_e$

arXiv:1512.08627

□ $M_{K\pi}$



$$BF(D^+ \rightarrow K^- \pi^+ e^+ \nu_e) = (3.71 \pm 0.03 \pm 0.08)\%$$

$$BF(D^+ \rightarrow K^- \pi^+ e^+ \nu_e)_{[0.8,1]} = (3.33 \pm 0.03 \pm 0.07)\%$$

□ Fractions of the component

$$f(D^+ \rightarrow (K^- \pi^+)_{K^{*0}(892)} e^+ \nu_e) = (93.93 \pm 0.22 \pm 0.18)\%$$

$$f(D^+ \rightarrow (K^- \pi^+)_{S\text{-wave}} e^+ \nu_e) = (6.05 \pm 0.22 \pm 0.18)\%$$

□ Parameters of $K^{*0}(892)$


$$m_{K^{*0}(892)} = (894.60 \pm 0.25 \pm 0.08) \text{ MeV}/c^2$$

$$\Gamma_{K^{*0}(892)} = (46.42 \pm 0.56 \pm 0.15) \text{ MeV}/c^2$$

Study of $D_s^+ \rightarrow \eta' X$ and $\eta' \rho^+$

PLB750 466(2015)

About 15.6 K ST D_s^- events by using 9 ST modes



$$\text{BF}(D_s^+ \rightarrow \eta' X) = (8.8 \pm 1.8 \pm 0.5)\%$$


Consistent with CLEO measurements $(11.7 \pm 1.8)\%$ [PRD79 112008(2009)]

$$\text{BF}(D_s^+ \rightarrow \eta' \rho^+) = (5.8 \pm 1.4 \pm 0.4)\%$$


Theoretical expectation $= (3.0 \pm 0.5)\%$ [PRD84 074019(2011)]

Resolve the disagreement between theoretical prediction and CLEO-c's previous measurement $\text{BF}(D_s^+ \rightarrow \eta' \rho^+) = (12.5 \pm 2.2)\%$ [PRD58 052002(1998)]

$$N(D_s^+ \rightarrow \eta' X) = 68 \pm 14$$



$$N(D_s^+ \rightarrow \eta' \rho^+) = 210 \pm 50$$



Other results

$D^+ \rightarrow \mu^+ \nu_\mu$ (Phys. Rev. D **89**, 051104(R) (2014))

$D^+ \rightarrow K_S^0 \pi^+ \pi^0$ (Phys. Rev. D **89**, 052001 (2014))

$D^0 \rightarrow \gamma\gamma$ and $D^0 \rightarrow \pi^0 \pi^0$ (Phys. Rev. D **91**, 112015 (2015))

Strong phase difference in $D^0 \rightarrow K^- \pi^+$ (Phys. Lett. B **734**, 227(2014))

y_{cp} in $D^0 - \bar{D}^0$ oscillation (Phys. Lett. B **744**, 339 (2015))

BF of D^{*0} decay (Phys. Rev. D **91**, 031101(R) (2015))

Observation of $\Lambda_c^+ \rightarrow n K_S^0 \pi^+$ BESIII preliminary

Summary

- With $D\bar{D}/D_s^+D_s^-/\Lambda_c^+\bar{\Lambda}_c^-$ produced at mass threshold, BESIII released many new results
 - Form factors measurement in (semi)leptonic decays of charmed hadrons provide important test to LQCD calculations and CKM matrix unitarity
 - Hadronic charmed hadrons decays improve the understanding of non-perturbative QCD
 - The first absolute BFs measurement of the Λ_c^+ hadronic decays
- BESIII is taking data at 4.18GeV to study physics related to D_s
- Many charm analyses are ongoing!

Thanks!